


Title: Quantum Mechanics from Gauge Theory

Date: Nov 24, 2004 03:50 PM

URL: <http://pirsa.org/04110039>


Abstract:

STRONG LOCALITY




$$\rho_{\lambda}^{AB}(x, y | \bar{i}, \bar{j}) = \rho_{\lambda}^A(x | i, \star) \rho_{\lambda}^B(y | \star, j)$$

STRONG LOCALITY




$$\rho_{\lambda}^{AB}(x, y | \bar{i}, \bar{j}) = \rho_{\lambda}^A(x | \bar{i}, *) \rho_{\lambda}^B(y | *, \bar{j})$$

STRONG LOCALITY




$$\rho_{\lambda}^{AB}(x, y | \bar{\epsilon}, \bar{j}) = \rho_{\lambda}^A(x | \epsilon, \star) \rho_{\lambda}^B(y | \star, \bar{j})$$

STRONG LOCALITY




$$\rho_{\lambda}^{AB}(x, y | \bar{i}, \bar{j}) = \rho_{\lambda}^A(x | \bar{i}, *) \rho_{\lambda}^B(y | *, \bar{j})$$

STRONG LOCALITY




$$\rho_{\lambda}^{AB}(x, y | \bar{\epsilon}, \bar{j}) = \rho_{\lambda}^A(x | \epsilon, \star) \rho_{\lambda}^B(y | \star, j)$$

STRONG LOCALITY




$$\rho_{\lambda}^{AB}(x, y | \bar{\epsilon}, \bar{j}) = \rho_{\lambda}^A(x | \epsilon, \star) \rho_{\lambda}^B(y | \star, j)$$

STRONG LOCALITY




$$\rho_{\lambda}^{AB}(x, y | \bar{\epsilon}, \bar{j}) = \rho_{\lambda}^A(x | \bar{\epsilon}, \star) \rho_{\lambda}^B(y | \star, \bar{j})$$

STRONG LOCALITY




$$\rho_{\lambda}^{AB}(x, y | \bar{\epsilon}, \bar{j}) = \rho_{\lambda}^A(x | \bar{\epsilon}, \star) \rho_{\lambda}^B(y | \star, \bar{j})$$

STRONG LOCALITY




$$\rho_{\lambda}^{AB}(x, y | \bar{\epsilon}, \bar{j}) = \rho_{\lambda}^A(x | \bar{\epsilon}, \star) \rho_{\lambda}^B(y | \star, \bar{j})$$

STRONG LOCALITY




$$\rho_{\lambda}^{AB}(x, y | \bar{\epsilon}, \bar{j}) = \rho_{\lambda}^A(x | \bar{\epsilon}, \star) \rho_{\lambda}^B(y | \star, \bar{j})$$

STRONG LOCALITY




$$\rho_{\lambda}^{AB}(x, y | \bar{\epsilon}, \bar{j}) = \rho_{\lambda}^A(x | \bar{\epsilon}, \star) \rho_{\lambda}^B(y | \star, \bar{j})$$

STRONG LOCALITY



$$\rho_{\lambda}^{AB}(x, y | \bar{\epsilon}, \bar{j}) = \rho_{\lambda}^A(x | \bar{\epsilon}, \bar{*}) \rho_{\lambda}^B(y | \bar{*}, \bar{j})$$

STRONG LOCALITY



$$\rho_{\lambda}^{AB}(x, y | \bar{i}, \bar{j}) = \rho_{\lambda}^A(x | \bar{i}, *) \rho_{\lambda}^B(y | *, \bar{j})$$